

Experimental test of the probability density function of true value of Poisson distribution parameter by single observation of number of events

S.I. Bityukov^{a*}, V.A. Medvedev^a, V.V. Smirnova^a, Yu.V. Zernii^b

^aInstitute for high energy physics, 142281 Protvino, Russia

^bMoscow State Academy of Instrument Engineering and Computer Science, Moscow, Russia

The empirical probability density function for the conditional distribution of the true value of Poisson distribution parameter on one measurement is constructed by computer experiment. The analysis of the obtained distributions confirms that these distributions are gamma-distributions.

1. INTRODUCTION

Let us consider the Gamma-distribution with probability density

$$g_x(\beta, \alpha) = \frac{x^{\alpha-1} e^{-\frac{x}{\beta}}}{\beta^\alpha \Gamma(\alpha)}. \quad (1)$$

At change of standard designations of Gamma-distribution $\frac{1}{\beta}$, α and x for a , $n+1$ and λ we get the following formula for probability density of Gamma-distribution

$$g_n(a, \lambda) = \frac{a^{n+1}}{\Gamma(n+1)} e^{-a\lambda} \lambda^n, \quad (2)$$

where a is a scale parameter and $n+1 > 0$ is a shape parameter. Suppose $a = 1$, then the probability density of Gamma-distribution $\Gamma_{1,n+1}$ looks like Poisson distribution of probabilities:

$$g_n(\lambda) = \frac{\lambda^n}{n!} e^{-\lambda}, \quad \lambda > 0, \quad n > -1. \quad (3)$$

Let the probability of observing n events in the experiment be described by a Poisson distribution with parameter λ , i.e.

$$f(n; \lambda) = \frac{\lambda^n}{n!} e^{-\lambda}. \quad (4)$$

*The participation in the Workshop is supported by the Organizing Committee of ACAT'03.

As it follows from the article [1] (see also [2]) and is clearly seen from the analysis of identity [3]

$$\sum_{k=n+1}^{\infty} f(k; \lambda_1) + \int_{\lambda_1}^{\lambda_2} g_n(\lambda) d\lambda + \sum_{k=0}^n f(k; \lambda_2) = 1, \quad (5)$$

i.e.

$$\sum_{k=n+1}^{\infty} \frac{\lambda_1^k e^{-\lambda_1}}{k!} + \int_{\lambda_1}^{\lambda_2} \frac{\lambda^n e^{-\lambda}}{n!} d\lambda + \sum_{k=0}^n \frac{\lambda_2^k e^{-\lambda_2}}{k!} = 1$$

for any $\lambda_1 \geq 0$ and $\lambda_2 \geq 0$, that at one measurement of the number of events n (in our case it is the number of casual events appearing in some system for certain period of time) which appear according to Poisson distribution, the parameter value of this distribution is described by Gamma-distribution $\Gamma_{1,1+n}$ with mean, mode, and variance $n+1$, n , and $n+1$, respectively. In other words conditional distribution of the probability of true value of parameter of Poisson distribution is a Gamma-distribution $\Gamma_{1,1+n}$ on condition that the measured value of the number of events is equal to n .

It means that the value n corresponds to the most probable value of parameter of Poisson distribution, and the mean value of the number of events, appearing in Poisson flow in the fixed time interval, must correspond to the magnitude $n+1$, i.e. the estimation of parameter of Poisson distribution at one observation is displaced for 1 from the measured value of the number of events. The

equation (5) in the considered case allows to mix Bayesian and frequentist probabilities.

As a result, we can easily construct the confidence intervals, take into account systematics and statistical uncertainties of measurements at statistical conclusions about the quality of planned experiments, estimate the value of the parameter of Poisson distribution by several observations [3,4].

Nevertheless there are works in which the approaches based on other assumptions of distribution of true value of parameter of Poisson distribution in presence of its estimation on a single measurement, for example [5]. Also the works using Monte Carlo methods for construction of confidence intervals and for estimations of Type I and Type II errors in the hypotheses testing have recently appeared [6,7] (see, also, [8]). Therefore the experimental test with the purpose to confirm, that the true value of parameter of Poisson distribution at single observation has density of probability of Gamma-distribution, and with the purpose to check up applicability of Monte Carlo methods to such tasks was carried out.

The structure of the paper is the following. In the next section the arrangement of computer experiment is considered, in the third section - the statistical analysis of the results is given, and the last section contains concluding remarks.

2. The arrangement of measurements

From the identity (5) follows, that any prior, except the uniform, on value of parameter of Poisson distribution in distribution of true value of this parameter at presence of the measured estimation n is excluded by existence of the boundary conditions determined by the appropriate sums of Poisson distributions (see Eq.(5)). Therefore we carried out the uniform scanning in parameter of Poisson distribution with step 0.1 from value $\lambda = 0.1$ up to value $\lambda = 20$, playing the Poisson distribution 30000 trials for each value λ (Fig.1) with the using of function RNPSSN [9].

After scanning for each value of number of the dropped out events n the empirical density of probability of true value of parameter of Poisson distribution to be λ if the observation is equal n




Figure 1. Amount of occurrences of n in the interval from 0 up to 9. Scanning in parameter of the Poisson distribution (30000 trials at each value of parameter λ) was carried out with step 0.1 in the interval of λ from 0.1 up to 20

was obtained.

3. The analysis of results

In Fig.2 the distribution (a), obtained at scanning (the playing of Poisson distribution with consecutive increase of the parameter λ after each series of trials with the fixed value of the parameter) in parameter λ with the selection of number of the dropped out events $n = 6$, and distribution (b) of the casual value, having $\Gamma_{1,7}$ distribution of appropriate area, calculated by the formula, are shown. One can see that the average value of parameter $\lambda \approx 7$. It means, the number of observed events is displaced for one unity by the estimation of the mean value of Poisson distribution parameter and correspond to the most probable value (the mean value has bias).

The same distributions obtained by the selection of number of dropped out events $n = 0$ (Fig.3) and $n = 8$ (Fig.4) superimposed on each other in logarithmic scale also are shown.

In Tab.1 the values of probabilities of compatibility of the empirical distribution, obtained by Monte Carlo by the scanning in parameter λ , and




Figure 2. Distributions of occurrences of value $n = 6$ depending on value of parameter λ . The distribution (a) is obtained at Monte Carlo scanning in parameter λ . The distribution (b) is obtained by direct construction of Gamma-distribution $\Gamma_{1,7}$.




Figure 3. Distributions of occurrences of value $n = 0$ depending on value of parameter λ . The distribution obtained at Monte Carlo scanning in parameter λ is superimposed on the distribution obtained by direct construction of Gamma-distribution $\Gamma_{1,1}$.




Figure 4. Distributions of occurrences of value $n = 8$ depending on value of parameter λ . The distribution obtained at Monte Carlo scanning in parameter λ is superimposed on the distribution obtained by direct construction of Gamma-distribution $\Gamma_{1,9}$.

the appropriate Gamma-distribution for values n from 0 up to 9 are presented. The calculations are based on the Kolmogorov Test (the function HDIFF of the package HBOOK [9]). The authors of a package as criterion of coincidence of two distributions recommend to use the requirement of probability value of compatibility more than 0.05. In Fig.4 the least continuous distributions are given.

Table 1
The probability of compatibility

n	probability	n	probability
0	1.000000	5	0.999084
1	0.999646	6	0.999986
2	0.992521	7	0.999892
3	0.999986	8	0.752075
4	0.999969	9	0.974236

Thus, the obtained results do not contradict the statement that conditional distribution of true value of parameter of Poisson distribution

at single measurement is obeyed to a Gamma-distribution.

4. Conclusion

In the report Monte Carlo experiment on the check of the statement, that true value of parameter of Poisson distribution at an estimation of this parameter on one observation n has probability density of Gamma-distribution $\Gamma_{1,n+1}$, is carried out. The obtained results confirm the conclusions of the papers [3,4] about a kind of conditional distribution of true value of parameter of Poisson distribution at single observation.

Note, that the given results also specify the applicability of Monte Carlo method for construction of conditional distribution of the true value of parameters of various distributions.

Acknowledgment

The authors thank N.V. Krasnikov, V.F. Obraztsov, V.A. Petukhov and M.N. Ukhanov for support of the given work. The authors also are grateful to S.S. Bitukov for fruitful discussions and constructive criticism. S.B. thank Toshiaki Kaneko and Fukuko Yuasa. The authors wish to thank E.A. Medvedeva for help in preparing the paper. This work has been supported by grant RFBR 03-02-16933.

REFERENCES

1. R.D. Cousins, Why isn't every physicist a Bayesian? *Am.J.Phys* **63** (1995) 398.
2. E.T. Jaynes: Papers on probability, statistics and statistical physics, Ed. by R.D. Rosenkrantz, D.Reidel Publishing Company, Dordrecht, Holland, 1983, p.165.
A.G.Frodesen, O.Skjeggestad, H.Toft, *Probability and Statistics in Particle Physics*, UNIVERSITETSFORLAGET, Bergen-Oslo-Tromso, 1979, p.408.
3. S.I. Bitukov, N.V. Krasnikov, V.A. Taperechkina, *Confidence intervals for Poisson distribution parameter*, Preprint IFVE 2000-61, Protvino, 2000; also, e-Print: hep-ex/0108020, 2001.
4. S.I.Bitukov, *On the Signal Significance in the Presence of Systematic and Statistical Uncertainties*, JHEP 09 (2002) 060, <http://www.iop.org/EJ/abstract/1126-6708/2002/09/06> Print: hep-ph/0207130.
S.I. Bitukov and N.V. Krasnikov, *Signal significance in the presence of systematic and statistical uncertainties*, Nucl.Instr.&Meth. **502** (2003) 795.
5. G.J. Feldman and R.D. Cousins, Unified approach to the classical statistical analysis of small signal, *Phys.Rev. D* **57** (1998) 3873-3889
6. S.I.Bitukov and N.V.Krasnikov, *On the observability of a signal above background*, Nucl.Instr.&Meth. **A452** (2000) 518.
7. J.Conrad et al., "Coverage of Confidence Intervals for Poisson Statistics in Presence of Systematic Uncertainties", Proc. of Conf. "Advanced statistical techniques in particle physics", eds. M.R. Whalley, L. Lyons, Durham, UK, 2002, p.58.
8. M.A. Stephens, *EDF statistics for goodness-of-fit and some comparison*, J. Amer.Statist.Assoc., **69**, 347 (1974) 730-737.
9. CERNLIB, CERN PROGRAM LIBRARY, Short Writeups, Entry V136 and Y250, (CERN, Geneva, Switzerland, Edition - June 1996)

Appendix

Let us consider the famous equality [1,2] in form as written in [3]

$$\sum_{i=n+1}^{\infty} f(i; \mu_1) + \int_{\mu_1}^{\mu_2} g(\mu; n) d\mu + \sum_{i=0}^n f(i; \mu_2) = 1, \quad (6)$$

$$\text{where } \mu_1 \geq 0, \mu_2 \geq 0, \\ f(k; \mu) = g(\mu; k) = \frac{\mu^k e^{-\mu}}{k!}, \quad k = 1, 2, \dots$$

Let us suppose that $g(\mu; k)$ is the probability density of parameter of the Poisson distribution to be μ if number of observed events is equal to k . It is a conditional probability density. As shown above (Eq.3) the $g(\mu; k)$ is the density of Gamma-distribution by definition.

On other side: if $g(\mu; k)$ is not equal to this probability density and the true probability density of the Poisson parameter is the other function $h(\mu; k)$ (continuous or with set of points of discontinuity by measure 0) then there takes place another identity

$$\sum_{i=n+1}^{\infty} f(i; \mu_1) + \int_{\mu_1}^{\mu_2} h(\mu; n) d\mu + \sum_{i=0}^n f(i; \mu_2) = 1, \quad (7)$$

This identity is correct for any $\mu_1 \geq 0$ and $\mu_2 \geq 0$.

If we subtract Eq.7 from Eq.6 then we have

$$\int_{\mu_1}^{\mu_2} (g(\mu; n) - h(\mu; n)) d\mu = 0. \quad (8)$$

We can choose the μ_1 and μ_2 by the arbitrary way. Let us make this choice so that $g(\mu; n)$ is not equal $h(\mu; n)$ in the interval (μ_1, μ_2) and, for example, $g(\mu; n) > h(\mu; n)$ and $\mu_2 > \mu_1$. In this case we have

$$\int_{\mu_1}^{\mu_2} (g(\mu; n) - h(\mu; n)) d\mu > 0 \quad (9)$$

and as a result we have contradiction. The identity (6) does not leave a place for any prior except uniform.